





Place for Rational & Educational Motivations!!

	$k = \frac{2.303}{t} \log \frac{[R]_0}{[R]}$	1
8	Henry's law states that the mole fraction of gas in the solution is proportional to the partial pressure of the gas over the solution. Applications: solubility of CO ₂ gas in soft drinks /solubility of air diluted with helium in blood used by sea divers or any other Solubility of gas in liquid decreases with increase in temperature.	1 ½ ½
9	X = CH ₃ -CO-CH ₂ -CH ₃ / Butan-2-one Y= CH ₃ -CH(OH)-CH ₂ -CH ₃ / Butan-2-ol	1 1
10	i)  ii) 	1+1
11	$k = \frac{2.303}{t} \log \frac{p_i}{2p_i - p_t}$ $= \frac{2.303}{300} \log \frac{0.3}{2 \times 0.3 - 0.5}$ $= \frac{2.303}{300} \log 3$ $= \frac{2.303 \times 0.4771}{300}$ $= 0.0036 \text{ atm}^{-1} \text{ or } 0.004 \text{ atm}^{-1} \text{ (approx.)}$	1 1 1




Place for Rational & Educational Motivations!!

12	i) Because of the resonance stabilization of the conjugate base i.e enolate anion or diagrammatic representation.	1½
	iii) Because the carboxyl group gets bonded to the catalyst anhyd. AlCl ₃ (Lewis acid). (note: part ii is deleted because of printing error and mark allotted in part i and part iii)	1½
	OR	
12	i) $C_6H_5CH_3 \xrightarrow{CrO_3/(CH_3CO)_2O} C_6H_5CH(OCOCH_3)_2 \xrightarrow{H_2O} C_6H_5CHO$ ii) $CH_3COOH \xrightarrow{Cl_2/P} Cl-CH_2-COOH$ iii) $CH_3COCH_3 \xrightarrow{Zn(Hg)/conc.HCl} CH_3CH_2CH_3$	1x3=3
	(Or by any other correct method)	
13	$d = \frac{z \times M}{N_A \times a^3}$ Or $d = \frac{z \times w}{N \times a^3}$ Where w is weight and N is no. of atoms. $d = \frac{4 \times 200 \text{ g}}{2.5 \times 10^{24} \times (400 \times 10^{-10} \text{ cm})^3}$ $d = 5 \text{ g cm}^{-3}$	1 1 1
	(or by any other correct method)	
14	i) It is a process in which both adsorption and absorption can take place simultaneously. ii) It is the potential difference between the fixed layer and the diffused/ double layer of opposite charges around the colloidal particles. iii) It is the temperature above which the formation of micelles takes place.	1 1 1







Place for Rational & Educational Motivations!!

15	$\Delta T_f = iK_f m$ <p>For complete ionisation of Na_2SO_4 $i=3$</p> $\Delta T_f = T_f^0 - T_f = 3 \times 1.86 \text{ K kg mol}^{-1} \times \frac{2\text{g}}{142\text{g mol}^{-1}} \times \frac{1000 \text{ g kg}^{-1}}{50 \text{ g}}$ $\Delta T_f = 1.57$ <p style="text-align: center;">So, $T_f = -1.57^\circ\text{C}$ or 271.43K</p>	<p style="text-align: right;">$\frac{1}{2}$</p> <p style="text-align: right;">$\frac{1}{2}$</p> <p style="text-align: right;">1</p> <p style="text-align: right;">1</p>
16	<p>i) Because of higher oxidation state (+5) / high charge to size ratio / high polarizing power.</p> <p>ii) Because of high interelectronic repulsion.</p> <p>iii) Because of its low bond dissociation enthalpy and high hydration enthalpy of F^-.</p>	<p>$1 \times 3 = 3$</p>
17	<p>i) A : $\text{C}_6\text{H}_5\text{CONH}_2$ B : $\text{C}_6\text{H}_5\text{NH}_2$ C : $\text{C}_6\text{H}_5\text{NHCOCH}_3$</p> <p>ii) A: $\text{C}_6\text{H}_5\text{NO}_2$ B : $\text{C}_6\text{H}_5\text{NH}_2$ C: $\text{C}_6\text{H}_5\text{-NC}$</p>	<p>$1\frac{1}{2}$</p> <p>$1\frac{1}{2}$</p>
18	<p>(i) Butadiene and acrylonitrile $\text{CH}_2 = \text{CH} - \text{CH} = \text{CH}_2$ and $\text{CH}_2 = \text{CH-CN}$</p> <p>(ii) Vinyl chloride $\text{CH}_2 = \text{CH-Cl}$</p> <p>(iii) Chloroprene</p> $\begin{array}{c} \text{Cl} \\ \\ \text{CH}_2 = \text{C} - \text{CH} = \text{CH}_2 \end{array}$	<p>$\frac{1}{2} + \frac{1}{2}$</p> <p>$\frac{1}{2} + \frac{1}{2}$</p> <p>$\frac{1}{2} + \frac{1}{2}$</p>
19	<p>i) </p> <p>ii) Peptide linkage / -CO-NH- linkage</p> <p>iii) Water soluble- Vitamin B / C Fat soluble- Vitamin A / D / E / K</p>	<p>1</p> <p>1</p> <p>$\frac{1}{2} + \frac{1}{2}$</p>





Place for Rational & Educational Motivations!!

20	<p>i) dsp^3, Diamagnetic, low spin</p> <p>ii) The energy used to split degenerate d-orbitals due to the presence of ligands in a definite geometry is called crystal field splitting energy.</p>	1 $\frac{1}{2} + \frac{1}{2}$ 1
21	<p>i) Iodine is heated with Zr or Ti to form a volatile compound which on further heating decompose to give pure Zr or Ti . or</p> $\text{Zr}(\text{impure}) + 2\text{I}_2 \longrightarrow \text{ZrI}_4 \text{ (volatile)}$ $\text{ZrI}_4 \xrightarrow{1800\text{K}} \text{Zr}(\text{pure}) + 2\text{I}_2$ <p>ii) Cryolite lowers the m.p. of alumina mix / acts as a solvent / brings conductivity.</p> <p>(iii) Role of NaCN in the extraction of Ag is to do the leaching of silver ore in the presence of air. or</p> $4\text{Ag}(\text{s}) + 8\text{CN}^-(\text{aq}) + 2\text{H}_2\text{O} + \text{O}_2(\text{g}) \longrightarrow 4[\text{Ag}(\text{CN})_2]^- + 4\text{OH}^-$	1 1 1
22	<p>i)</p>  <p>ii)</p>  <p>iii) $\text{CH}_3\text{CH}_2\text{ONO}$</p>	1 x 3 = 3





Place for Rational & Educational Motivations!!

23	<p>(i)Caring ,dutiful, Concerned, compassionate (or any other two values)</p> <p>ii)Because higher doses may have harmful effects and act as poison which cause even death.</p> <p>iii)Tranquilizers are a class of chemical compounds used for treatment of stress or even mental diseases. ex. chlordiazepoxide, equanil,veronal, serotonin,valium (or any other two examples)</p>	<p>½+½</p> <p>1</p> <p>1</p> <p>½+½</p>
24	<p>a)</p> <p>Given $E^{\circ}_{\text{Cell}} = + 0.30\text{V}$; $F = 96500\text{C mol}^{-1}$</p> <p>$n = 6$ (from the given reaction)</p> <p>$\Delta_r G^{\circ} = - n \times F \times E^{\circ}_{\text{Cell}}$</p> <p>$\Delta_r G^{\circ} = - 6 \times 96500 \text{ C mol}^{-1} \times 0.30\text{V}$</p> <p>$= - 173,700 \text{ J / mol or } - 173.7 \text{ kJ / mol}$</p> <p>$\log K_c = \frac{n E^{\circ}_{\text{Cell}}}{0.059}$</p> <p>$\log K_c = \frac{6 \times 0.30}{0.059}$</p> <p>$\log K_c = 30.5$</p> <p>b)A Because E° value of A shows that on coating ,A acts as anode and Fe acts as a cathode and hence A oxidises in preference to Fe and prevent corrosion / or E°_{cell} is positive and hence A oxidises itself to prevent corrosion of Fe/E° value is more negative. (or any other correct reason)</p> <p align="center">OR</p>	<p>½</p> <p>1</p> <p>½</p> <p>1</p> <p>1</p> <p>1</p>



Place for Rational & Educational Motivations!!

24	<p>a)</p> $\Lambda_m = \frac{\kappa}{c}$ $= \frac{3.905 \times 10^{-5} \text{ S cm}^{-1}}{0.001 \text{ mol L}^{-1}} \times \frac{1000 \text{ cm}^3}{\text{L}}$ $\Lambda_m = 39.05 \text{ Scm}^2 \text{ mol}^{-1}$ $\Lambda_0 = \lambda^0(\text{H}^+) + \lambda^0(\text{CH}_3\text{COO}^-)$ $= (349.6 + 40.9) \text{ Scm}^2 \text{ mol}^{-1}$ $\Lambda_0 = 390.5 \text{ Scm}^2 \text{ mol}^{-1}$ $\alpha = \frac{\Lambda_m}{\Lambda_0}$ $= \frac{39.05 \text{ Scm}^2 \text{ mol}^{-1}}{390.5 \text{ Scm}^2 \text{ mol}^{-1}}$ $\alpha = 0.1$ <p>b) Secondary battery or rechargeable battery</p> $\text{Pb(s)} + \text{PbO}_2(\text{s}) + 2\text{SO}_4^{2-}(\text{aq}) + 4\text{H}^+(\text{aq}) \longrightarrow 2\text{PbSO}_4(\text{s}) + 2\text{H}_2\text{O}(\text{l})$	<p>1/2</p> <p>1</p> <p>1/2</p> <p>1</p> <p>1</p>
25	<p>a)</p> <p>i) Because of higher oxidation state (+7) of Mn.</p> <p>ii) Because it has one unpaired electron in 3d orbital in its +2 oxidation state / or it has incompletely filled d-orbital in +2 oxidation state.</p> <p>iii) Because of comparable energies of 5f, 6d and 7s orbitals.</p> <p>b)</p> $2\text{MnO}_2 + 4\text{KOH} + \text{O}_2 \longrightarrow 2\text{K}_2\text{MnO}_4 + 2\text{H}_2\text{O}$ $3\text{MnO}_4^{2-} + 4\text{H}^+ \longrightarrow 2\text{MnO}_4^- + \text{MnO}_2 + 2\text{H}_2\text{O}$	<p>1</p> <p>1</p> <p>1</p> <p>1+1</p>
OR		

26	<p>a). (i)</p>  <p>(ii) $\text{CH}_3\text{CH}_2\text{OH} + \text{CH}_3\text{COCl} \xrightarrow{\text{pyridine}} \text{CH}_3\text{CH}_2\text{O-COCH}_3 + \text{HCl}$</p> <p>(iii).</p>  <p>(b)(i) Warm each compound with iodine and sodium hydroxide. Phenol : No yellow ppt formed Ethanol: Yellow ppt of Iodoform are formed.</p> <p>ii) On adding lucas reagent ($\text{HCl}/\text{anhyd. ZnCl}_2$), Propan-2-ol gives white turbidity after 5 minutes whereas 2-methylpropan-2-ol gives white turbidity immediately. (or any other suitable test)</p>	<p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p>
----	---	--